Technical support: support@abbkine.com

Website: https://www.abbkine.com

GSK3α/β (phospho Tyr279/216) Polyclonal Antibody

Cat #: ABP50322 Size: 30µl /100µl /200µl

Product Information

	Product Name: GSK3α/β (phospho Tyr279/216) Polyclonal Antibody		
	Applications: WB, IF, IHC-P, ELISA		Isotype: Rabbit IgG
	Reactivity: Human, Mouse, Rat		
REF	Catalog Number: ABP50322	LOT	Lot Number: Refer to product label
	Formulation: Liquid		Concentration: 1 mg/ml
Ĵ.	Storage: Store at -20°C. Avoid repeated freeze / thaw cycles.	<u>∧</u>	Note: Contain sodium azide.

Background: GSK3A encodes a multifunctional Ser/Thr protein kinase (glycogen synthase kinase 3 alpha) that is implicated in the control of several regulatory proteins including glycogen synthase, and transcription factors, such as JUN. It also plays a role in the WNT and PI3K signaling pathways, as well as regulates the production of beta-amyloid peptides associated with Alzheimer's disease.

<u>Application Notes</u>: Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows: WB (1:500-1:2000), IHC-P (1:100-1:300), ELISA (1:20000). Not yet tested in other applications.

Storage Buffer: PBS containing 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.

Storage Instructions: Stable for one year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.




Fig.1. Immunohistochemical analysis of paraffinembedded human tonsil tissue. 1, GSK3 α / β (phospho Tyr279/216) Polyclonal Antibody was diluted at 1:200 (4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C, 20min). 3, secondary






Fig.2. Immunohistochemical analysis of paraffinembedded mouse heart tissue. 1, GSK3 α / β (phospho Tyr279/216) Polyclonal Antibody was diluted at 1:200 (4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C, 20min). 3, secondary antibody was diluted at 1:200 (room temperature, 30min). Negative control was used by secondary antibody only.

Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

