


Phospho-Caspase 8 (Tyr448) Ab

Cat.#: AF8103
 Size: 100ul,200ul

Concn.: 1mg/ml
 Source: Rabbit


Mol.Wt.: 57kDa
 Clonality: Polyclonal

- Application:** WB 1:1000-3000, IHC 1:50-1:200, IF/ICC 1:100-1:500
- Reactivity:** Human,Mouse,Rat
- Purification:** The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
- Specificity:** Caspase 8 (Phospho-Tyr448) Ab detects endogenous levels of Caspase 8 only when phosphorylated at Tyr448.
- Immunogen:** A synthesized peptide derived from human Caspase 8 (Phospho-Tyr448).
- Uniprot:** Q14790
- Subcellular Location:** Cytoplasm.
- Tissue Specificity:** Isoform 1, isoform 5 and isoform 7 are expressed in a wide variety of tissues. Highest expression in peripheral blood leukocytes, spleen, thymus and liver. Barely detectable in brain, testis and skeletal muscle.
- Similarity:** Isoform 9 contains a N-terminal extension that is required for interaction with the BCAP31 complex.Belongs to the peptidase C14A family.
- Storage Condition and Buffer:** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.




Western blot analysis of Caspase 8 (Phospho-Tyr448) using forskolin treated HeLa whole cell lysates.


-/+ means absence or presence of N peptide (non-phospho peptide) and P peptide (phospho peptide).




AF8103 at 1/200 staining Rat spleen tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



AF8103 at 1/200 staining Mouse kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



AF8103 at 1/200 staining Human kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



AF8103 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.