

CXCL17 Ab

Cat.#: DF10167
Size: 100ul,200ul


Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 14 kDa
Clonality: Polyclonal

Application:	WB 1:1000-3000
Reactivity:	Human
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).
Specificity:	CXCL17 Ab detects endogenous levels of total CXCL17.
Immunogen:	A synthesized peptide derived from human CXCL17.
Uniprot:	Q6UXB2
Subcellular Location:	Extracellular region or secreted;
Tissue Specificity:	Detected in trachea, stomach, lung and skeletal muscle. Detected in intestine and in normal and asthmatic lung (at protein level). Breast tumors showed 3- to 24-fold up-regulation.
Similarity:	Belongs to the intercrine alpha (chemokine CxC) family.
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from MCF7, using CXCL17 Ab. The lane on the left was treated with blocking peptide.



Western blot analysis of CXCL17 using HUVEC whole cell lysates

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.