Phospho-EGFR (Tyr1068) Ab

Cat.#: DF2969 Concn.: 1mg/ml Mol.Wt.: 180 KD Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:1000-3000, IF/ICC 1:100-1:500

Reactivity: Human, Mouse, Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-EGFR (Tyr1068) Ab detects endogenous levels of

EGFR only when phosphorylated at Tyr1068.

Immunogen: A synthesized peptide derived from human Phospho-EGFR (

Tyr1068).

Uniprot: P00533

Subcellular Location: Secreted and Cell membrane. Endoplasmic reticulum

membrane. Golgi apparatus membrane. Nucleus membrane. Endosome. Endosome membrane. Nucleus. In response to EGF, translocated from the cell membrane to the nucleus via Golgi and ER. Endocytosed upon activation by ligand. Colocalized with GPER1 in the nucleus of estrogen agonist-

induced cancer-associated fibroblasts (CAF).

Tissue Specificity: Ubiquitously expressed. Isoform 2 is also expressed in

ovarian cancers.

Similarity: Belongs to the protein kinase superfamily. Tyr protein kinase

family. EGF receptor subfamily.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.

kDa 250— 150— 100— 75— 50— 37— 25— 20—

Western blot analysis of Phospho-EGFR (Tyr1068) using

COLO205 whole cell lysates



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



DF2969 staining SK-OV3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.