


Phospho-Nephrin(Ser432) Ab

Cat.#: AF7452
Size: 100ul,200ul


Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 135 kDa
Clonality: Polyclonal

- Application:** WB 1:500-1:2000, IHC 1:50-1:200
- Reactivity:** Human,Mouse
- Purification:** The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
- Specificity:** Phospho-Nephrin(Ser432) Ab detects endogenous levels of Nephrin only when phosphorylated at Ser432.
- Immunogen:** A synthesized peptide derived from human Nephrin around the phosphorylation site of Ser432.
- Uniprot:** O60500
- Subcellular Location:** Cell membrane. Predominantly located at podocyte slit diaphragm between podocyte foot processes. Also associated with podocyte apical plasma membrane.
- Tissue Specificity:** Specifically expressed in podocytes of kidney glomeruli.
- Similarity:** Belongs to the immunoglobulin superfamily.
- Storage Condition and Buffer:** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from mouse kidney tissue, using Phospho-Nephrin(Ser432) Ab. Lane 1 was treated with phospho-blocking peptide, Lane 2 was treated with non-phospho-blocking peptide.



Western blot analysis of extracts from mouse kidney tissue sample, using Phospho-Nephrin(Ser432) Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.