

Phospho-Atg14 (Ser29) Ab

Cat.#: AF2320
Size: 100ul,200ul

Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 65kDa
Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200

Reactivity: Human,Mouse

Purification: The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Specificity: Phospho-Atg14 (Ser29) Ab detects endogenous levels of Atg14.


Immunogen: A synthesized peptide derived from human Atg14 around the phosphorylation site of Ser29.

Uniprot: Q6ZNE5


Subcellular Location: Cytoplasm. Endoplasmic reticulum. Cytosolic under nutrient-rich conditions. Following autophagy stimuli, such as starvation or rapamycin induction, predominantly detected in cytoplasmic foci, identified as isolation membranes and autophagosomes.

Similarity: The coiled-coil domain is required for BECN1- and PIK3C3-binding and for autophagy.The final 80 residues in the C-terminus define a minimum required region for autophagosome binding called BATS.The N-terminal cysteine repeats are required for proper localization to the endoplasmic reticulum.Belongs to the ATG14 family.


Storage Condition and Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from A172 cells, using Phospho-Atg14 (Ser29) Ab. Lane1 was treated with phospho-blocking peptide, Lane2 was treated with non-phospho-blocking peptide.



Western blot analysis of extracts from A172 cells, using Phospho-Atg14 (Ser29) Ab.



AF2320 at 1/100 staining Human thyroid cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.