


Phospho-Cortactin (Tyr421) Ab

Cat.#: AF3436 Size: 100ul,200ul	Concn.: 1mg/ml Source: Rabbit	Mol.Wt.: 85kDa Clonality: Polyclonal
Application:	WB 1:500-1:2000, IF/ICC 1:100-1:500	
Reactivity:	Human,Mouse,Rat	
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.	
Specificity:	Phospho-Cortactin (Tyr421) Ab detects endogenous levels of Cortactin only when phosphorylated at Tyrosine 421.	
Immunogen:	A synthesized peptide derived from human Cortactin around the phosphorylation site of Tyrosine 421.	
Uniprot:	Q14247	
Description:	cortactin a cytoskeletal protein coordinating actin reorganizatio Localizes at the leading edge of migration. Its amino-terminal ac the Arp2/3 and WASP complex a	n during cell movement. Flamellipodia during cell cidic domain associates with
Subcellular Location:	Cytoplasm > cytoskeleton. Cell Cell projection > ruffle. Associat and lamellipodia.	
Similarity:	The SH3 motif may mediate binding to the cytoskeleton.	
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.	



Western blot analysis of Phospho-Cortactin (Tyr421) expression in various lysates





AF3436 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.