


Phospho-HDAC2 (Ser394) Ab

Cat.#: AF3470
 Size: 100ul,200ul


Concn.: 1mg/ml
 Source: Rabbit

Mol.Wt.: 55kDa
 Clonality: Polyclonal

Application:	WB 1:500-1:2000, IF/ICC 1:100-1:500
Reactivity:	Human,Mouse,Rat
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Specificity:	Phospho-HDAC2 (Ser394) Ab detects endogenous levels of HDAC2 only when phosphorylated at Serine 394.
Immunogen:	A synthesized peptide derived from human HDAC2 around the phosphorylation site of Serine 394.
Uniprot:	Q92769
Description:	HDAC2 a transcriptional regulator of the histone deacetylase family, subfamily 1. Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation plays a role in epigenetic repression and transcriptional regulation, cell cycle progression and developmental events.
Subcellular Location:	Nucleus.
Tissue Specificity:	Widely expressed; lower levels in brain and lung.
Similarity:	Belongs to the histone deacetylase family. HD type 1 subfamily.
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from rat spleen, using Phospho-HDAC2 (Ser394) Ab.



Western blot analysis of Phospho-HDAC2 (Ser394) expression in various lysates



AF3470 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab (Red), diluted at 1/600, was used as secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.