

## G3BP-1 Ab

Cat.#: AF6427 Concn.: 1mg/ml Mol.Wt.: 60kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000 IHC 1:50-1:200

Reactivity: Human, Mouse

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: G3BP-1 Ab detects endogenous levels of total G3BP-1.

Immunogen: A synthesized peptide derived from human G3BP-1.

Uniprot: Q13283

Description: G3BP-1 an hnRNA-binding protein and endoribonuclease that

participates in the Ras signal transduction pathway. A regulated effector of stress granule (SG) assembly. SGs are involved in mRNA sorting in the storing of untranslated

mRNAs.

Subcellular Location: Cytoplasm > cytosol. Cell membrane. Nucleus.

Cytoplasmic in proliferating cells, can be recruited to the plasma membrane in exponentially growing cells (By similarity). Cytosolic and partially nuclear in resting cells. Recruited to stress granules (SGs) upon either arsenite or high temperature treatment. Recruitment to SGs is

influenced by HRAS.

Tissue Specificity: Ubiquitous.

Similarity: The NTF2 domain mediates multimerization.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.

280 — 180 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 — 190 —

Western blot analysis of G3BP-1 expression in 293 whole cell

lysates



## Affinity Biosciences

website:www.affbiotech.com order:order@affbiotech.com



AF6427 at 1/100 staining Mouse colon tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.