

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

c-MYC Ab

Cat.#: AF0358 Concn.: 1mg/ml Mol.Wt.: 49kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB: 1:500~1:3000 IHC: 1:50~1:200, IF/ICC 1:100-1:500

Reactivity: Human, Mouse, Rat

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: MYC Ab detects endogenous levels of total MYC.

Immunogen: A synthesized peptide derived from human MYC.

Uniprot: P01106

Description: Myc a proto-oncogenic transcription factor that plays a role

in cell proliferation, apoptosis and in the development of human tumors.. Seems to activate the transcription of

growth-related genes.


Subcellular Location: Nucleus > nucleoplasm. Nucleus > nucleolus.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20

°C.Stable for 12 months from date of receipt.



Western blot analysis on 293 cell lysates using MYC Ab, The lane on the left was treated with the antigen-specific peptide.



AF0358 at 1/100 staining Human breast cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



AF0358 staining 293 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween020 at 4° C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.