

MBL2 Ab

Cat.#: DF4152	Concn.: 1mg/ml	Mol.Wt.: 27 KD
Size: 100ul,200ul	Source: Rabbit	Clonality: Polyclonal

Application: WB 1:500~1:1000, IF/ICC 1:100-1:500

Reactivity: Human,Mouse,Rat

Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Specificity: MBL2 Ab detects endogenous levels of total MBL2.

Immunogen: A synthesized peptide.


Uniprot: P11226

Subcellular Location: Secreted.


Tissue Specificity: Plasma protein produced mainly in the liver.

Similarity: The coiled-coil domain mediates trimerization.

Storage Condition and Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of MBL2 expression in various lysates



DF4152 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking,



Affinity Biosciences
website: www.affbiotech.com
order: order@affbiotech.com

overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.