

SOX2 Ab

Cat.#: DF6109
Size: 100ul, 200ul

Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 34kDa
Clonality: Polyclonal

Application: WB 1:500-1:2000 IHC 1:50-1:200

Reactivity: Human, Mouse, Rat

Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Specificity: SOX2 Ab detects endogenous levels of total SOX2.


Immunogen: A synthesized peptide derived from human SOX2.

Uniprot: P48431

Description: Embryonic stem cells are derived from the inner cell mass of the blastocyst and are unique in their pluripotent capacity and potential for self-renewal. Sox2 is one of a set of transcription factors that are crucial for the maintenance of pluripotency (1). Sox2, Oct-4, and Nanog cooperate in this network (1-3), and siRNA knockdown of either Sox2 or Oct-4 results in loss of pluripotency (4,5). Chromatin immunoprecipitation experiments have shown that Sox2 and Oct-4 bind to thousands of gene regulatory sites, highlighting the importance of these transcription factors in early embryonic development (6,7). It has recently been shown that Sox2 is amplified in lung and esophageal squamous cell tumors (8).

Subcellular Location: Nucleus.

Storage Condition and Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from date of receipt.



Western blot analysis of human brain lysates using SOX2 Ab. The lane on the left was treated with the antigen-specific peptide.



DF6109 at 1/100 staining Mouse brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.