

## GATA2 Ab

Cat.#: DF6117  
Size: 100ul,200ul

Concn.: 1mg/ml  
Source: Rabbit

Mol.Wt.: 49kDa  
Clonality: Polyclonal

Application: WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500

Reactivity: Human,Mouse,Rat

Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Specificity: GATA2 Ab detects endogenous levels of total GATA2.

Immunogen: A synthesized peptide derived from human GATA2.


Uniprot: P23769

Description: The zinc finger transcription factor GATA-2 is widely expressed and plays an essential role in many developmental processes (1). Studies on GATA-2 knockout mice indicate that this protein is required in hematopoiesis (2). GATA-2 also inhibits the differentiation of white (3) and brown adipocytes (4) and has been shown to suppress the proliferation of neuronal progenitor cells (5).


Subcellular Location: Nucleus.

Tissue Specificity: Endothelial cells.

Storage Condition and Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of Jurkat lysates using GATA2 Ab. The lane on the left was treated with the antigen-specific peptide.



DF6117 at 1/100 staining Mouse liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



DF6117 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.