


Histone H4K20me3 Ab

Cat.#: DF6950 Size: 100ul,200ul	Concn.: 1mg/ml Source: Rabbit	Mol.Wt.: 11kDa Clonality: Polyclonal
Application:	WB 1:500-1:1000	
Reactivity:	Human,Mouse	
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).	
Specificity:	Histone H4K20me3 Ab detects endogenous levels of total Histone H4K20me3.	
lmmunogen:	A synthetic methylated peptide corresponding to residues surrounding K20 of human histone H4.	
Uniprot:	P62805	
Description:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the centromeric copy.	
Subcellular Location:	Nucleus. Chromosome.	
Similarity:	Belongs to the histone H4 family.	
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.	





Western blot analysis of Hela whole cell lysates, using Histone H4K20me3 Ab. The lane on the left was treated with the antigen-specific peptide.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.