


BIK Ab

Cat.#: DF6958
Size: 100ul,200ul


Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 18kDa
Clonality: Polyclonal

Application:	WB 1:500-1:2000 IHC 1:50-1:200, IF/ICC 1:100-1:500
Reactivity:	Human
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).
Specificity:	BIK Ab detects endogenous levels of total BIK.
Immunogen:	A synthesized peptide derived from human BIK.
Uniprot:	Q13323
Description:	Bik/Nbk (Bcl-2-interacting killer/natural born killer) is a potent pro-apoptotic protein belonging to a group of Bcl-2 family members that includes Bad, Bid, Bim, Hrk, and Noxa, containing a BH3 domain but lacking other conserved domains, BH1 or BH2 (1,2). Functionally, Bik is able to bind to and antagonize anti-apoptotic Bcl-2 family members including Bcl-2, Bcl-xL, and viral homologs E1B-19K and EBV-BHFR1. The BH3 domain of Bik is essential for its apoptotic activity and interaction with survival proteins (3). Phosphorylation of Bik is correlated with an increase in its pro-apoptotic activity (4).
Subcellular Location:	Endomembrane system. Around the nuclear envelope, and in cytoplasmic membranes.
Similarity:	Intact BH3 motif is required by BIK, BID, BAK, BAD and BAX for their pro-apoptotic activity and for their interaction with anti-apoptotic members of the Bcl-2 family.
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from Raji , using BIK Ab. The lane on the left was treated with the antigen-specific peptide.



DF6958 at 1/100 staining Human gastric tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



DF6958 staining HepG2 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.