

PAB874Mu02

Polyclonal Antibody to Coagulation Factor VII (F7)
Organism Species: Mus musculus (Mouse)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



### [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific Affinity Chromatography.

Traits: Liquid

Concentration: 200µg/mL

**UOM**: 100µg

Applications: WB; IHC; ICC; IP.

### [ IMMUNOGEN ]

Immunogen: Recombinant F7 (Ala42~Lys169) expressed in E.coli.

Accession No.: RPB874Mu02

## [APPLICATIONS]

Western blotting: 0.5-5µg/mL

Immunohistochemistry: 5-20µg/mL Immunocytochemistry: 5-20µg/mL

Optimal working dilutions must be determined by end user.

### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

## [ QUALITY CONTROL ]

Content: The quality control contains recombinant F7 disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol, 0.02% BPB, 50mM DTT and 0.02% NaN<sub>3</sub>.

## [STORAGE AND STABILITY]


Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]



#### Figure 1. Western Blot

A. Sample: Recombinant F7, Mouse

B. Lane1: Mouse Placenta Tissue

Lane2: Mouse Spleen Tissue

Primary Ab: 3µg/mL Rabbit Anti-Mouse F7 Ab

Second Ab: 1:2000 Dilution of HRP-Linked Guinea pig

Anti-Rabbit Ab (Catalog: SAA544Rb59)




Figure 2. DAB staining on IHC-P

Samples:

A. Mouse Liver Tissue

B. Mouse Kidney Tissue