

RPD178Hu01 10µg
Recombinant Paraoxonase 3 (PON3)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[**PROPERTIES**]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Gly2~Leu354

Tags: N-terminal His-Tag

Tissue Specificity: Liver.

Subcellular Location: Secreted.

Purity: >98%

Traits: Freeze-dried powder

Buffer formulation: 100mM NaHCO₃, 500mM NaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.2

Predicted Molecular Mass: 43.2kDa

Accurate Molecular Mass: 43kDa as determined by SDS-PAGE reducing conditions.

[**USAGE**]

Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```

GKLVALVLL  GVGLSLVGEM  FLAFRERVNA  SREVEPVEPE  NCHLIEELES
GSEDIDILPS  GLAFISSGLK  YPGMPNFAPD  EPGKIFLMDL  NEQNPRQAAL
EISGGFDKEL  FNPHGISIFI  DKDNTVYLYV  VNHPHMKSTV  EIFKFEEQQR
SLVYLKTIKH  ELLKSVNDIV  VLGPEQFYAT  RDHYFTNSLL  SFFEMILDLR
WTYVLFYSPR  EVKVVAKGFC  SANGITVSAD  QKYVYVADVA  AKNIHIMEKH
DNWDLTQLKV  IQLGTLVDNL  TVDPATGDIL  AGCHPNPMKL  LNYNPEDPPG
SEVLRIQNVL  SEKPRVSTVY  ANNGSVLQGT  SVASVYHGKI  LIGTVFHKTL
YCEL
    
```

[IDENTIFICATION]




Figure 1. SDS-PAGE