



## **Beta Lactamase Mixture Supplement w/ 2K IU of Cephalosporinase and 20K IU of Penicillinase**

**FD289**

An innovative enzyme based product that can efficiently inactivate wide range of antibiotics like Penicillins, Cephalosporins of first, second, third and fourth generation and Penems.

### **Composition** (per vial contains)

### **Ingredients**

Cephalosporinase	> 2000 IU activity
Penicillinase	>20000 IU activity

1 IU is defined as the amount of enzyme needed to hydrolyze 1 $\mu$ mole of Penicillin G (Penicillinase) or 1 $\mu$ mole of Cephalosporin C (Cephalosporinase) per minute at 25°C and pH 7.0.

1 IU of Penicillinase corresponds to 600 Levy Units or 75 Pollock Units.

### **Directions:**

Rehydrate the contents of 1 vial with 20 ml of distilled water. After reconstitution, filter sterilize the solution with 0.2 micron syringe filter. Add appropriate amount of solution depending on the application. Remaining sterilized solution can be stored at 2-8°C for 4 weeks.

### **Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.