

Product	Code	Packing
L-Tyrosine disodium salt, dihydrate	TC477-25G	25gm
L-3-(4-Hydroxyphenyl)alanine disodium salt (From non-animal source)	TC477-100G	100gm
Cell Culture Tested $C_9H_9NO_3 \cdot 2Na \cdot 2H_2O$ MW : 261.19 CAS : 122666-87-9 Assay : $\geq 95\%$ Store below 30°C	TC477-500G	500gm
L-Valine	TC132-25G	25gm
(From non-animal source)	TC132-100G	100gm
(L-2-Amino-3-methylbutanoic acid)	TC132-500G	500gm
Cell Culture Tested $C_6H_{11}NO_2$ MW : 117.15 CAS : 72-18-4 Assay : $\geq 99\%$ Store below 30°C	TC132-1KG	1kg

Antibiotics

Contamination Control Antibiotics

Actinomycin D	TC018-5MG	5mg
Cell Culture Tested	TC018-2X5MG	2x5mg
Recommended for use in cell culture applications at 1mg/L	TC018-5X5MG	5x5mg
$C_{62}H_{86}N_{12}O_{16}$ MW : 1255.42 CAS : 50-76-0 Assay : $\geq 97.5\%$ Store at 2 - 8°C		
Amphotericin B	TC019-1G	1gm
Cell Culture Tested	TC019-5G	5gm
Recommended for use in cell culture applications at 2.5 mg/L		
$C_{47}H_{73}NO_{17}$ MW : 924.08 CAS : 1397-89-3 Assay : $\geq 90\%$ Potency : $\geq 750 \mu\text{g}/\text{mg}$ Store at 2 - 8°C		
Ampicillin sodium salt	TC021-1G	1gm
Cell Culture Tested	TC021-5G	5gm
Recommended for use in cell culture applications at 100 mg/L	TC021-25G	25gm
$C_{16}H_{18}N_3NaO_4S$ MW : 371.39 CAS : 69-52-3 Assay : $\geq 91\%$ Potency : $\geq 845 \mu\text{g}/\text{mg}$ Store at 2 - 8°C		
Bacitracin	TC201-1VL	1vl
50,000 units per vial	TC201-5VL	5vl
Cell Culture Tested $C_{66}H_{103}N_{17}O_{16}S$ MW : 1422.69 CAS : 1405-87-4 Potency : $\geq 60 \text{ units}/\text{mg}$ Store at 2 - 8°C		
*Blasticidin S Hydrochloride	TC421-25MG	25mg
Blasticidin S	TC421-100MG	100mg
Cell Culture Tested $C_{17}H_{26}N_8O_5 \cdot HCl$ MW : 458.9 CAS : 3513-03-9 Assay : $\geq 99.5\%$ Store at 2 - 8°C	TC421-1G	1gm

Product	Code	Packing
Carbenicillin disodium salt	TC199-250MG	250mg
Cell Culture Tested	TC199-1G	1gm
$C_{17}H_{18}N_2O_5SNa_2$	TC199-5G	5gm
MW : 422.36 CAS : 4800-94-6	TC199-10G	10gm
Assay : $\geq 90\%$ Potency : $\geq 770 \mu\text{g}/\text{mg}$ Store at 2 - 8°C		
Cephalothin sodium salt	TC022-250MG	250mg
Cell Culture Tested	TC022-1G	1gm
Recommended for use in cell culture applications at 100 mg/L	TC022-5G	5gm
$C_{16}H_{15}N_3NaO_5S_2$ MW : 418.42 CAS : 58-71-9 Assay : $\geq 95\%$ Store at 2 - 8°C		
*Cefotaxime Sodium Salt	TC352-1G	1gm
Cell Culture Tested	TC352-5G	5gm
$C_{16}H_{15}N_5O_5S_2Na$	TC352-25G	25gm
MW : 477.45 CAS : 64485-93-4		
Assay : $\geq 98\%$ Potency : 916-964 $\mu\text{g}/\text{mg}$ Store at 2 - 8°C		
Chloramphenicol	TC204-5G	5gm
Cell Culture Tested	TC204-25G	25gm
$C_{11}H_{12}Cl_2N_2O_5$ MW : 323.13 CAS : 56-75-7 Assay : $\geq 98\%$ Store below 30°C		
Ciprofloxacin hydrochloride monohydrate	TC447-1G	1gm
Cell Culture Tested	TC447-5G	5gm
$C_{17}H_{18}FN_3O_3 \cdot HCl \cdot H_2O$	TC447-25G	25gm
MW : 385.82 CAS : 86393-32-0 Assay : $\geq 98\%$ Store below 30°C		
Cycloheximide	TC196-1G	1gm
Cell Culture Tested	TC196-5G	5gm
$C_{15}H_{23}NO_4$ MW : 281.35 CAS : 66-81-9 Assay : $\geq 96\%$ Store at 2 - 8°C		
Erythromycin	TC024-1G	1gm
Cell Culture Tested	TC024-5G	5gm
Recommended for use in cell culture applications at 100 mg /L		
$C_{37}H_{67}NO_{13}$ MW : 733.93 CAS : 114-07-8 Assay : $\geq 98\%$ Potency : $\geq 920 \mu\text{g}/\text{mg}$ Store below 30°C		
Gentamicin sulphate	TC026-1G	1gm
Cell Culture Tested	TC026-5G	5gm
Recommended for use in Cell Culture application at 50 mg/L	TC026-25G	25gm
$CAS : 1405-41-0$ Potency : $\geq 590 \text{ IU}/\text{mg}$ Store at 2 - 8°C		

* Delivery after 4 - 6 weeks

Unless indicated all the above products to be stored below 30°C.