

Product	Code	Packing
Spectinomycin dihydrochloride pentahydrate Cell Culture Tested Recommended for use in cell culture application at 7.5-20 mg/L $C_{14}H_{24}N_2O_7 \cdot 2HCl \cdot 5H_2O$ MW : 495.35 CAS : 22189-32-8 Potency : $\geq 603 \mu\text{g}/\text{mg}$ Store at 2 - 8°C	TC034-1G TC034-5G TC034-25G	1gm 5gm 25gm
Streptomycin sulphate Cell Culture Tested Recommended for use in cell culture application at 100 mg/L $C_{21}H_{39}N_7O_{12} \cdot 1.5 H_2SO_4$ MW : 728.69 CAS : 3810-74-0 Potency : $\geq 730 \text{ Units}/\text{mg}$ Store at 2 - 8°C	TC035-5G TC035-25G	5gm 25gm
Tetracycline hydrochloride Cell Culture Tested Recommended for use in Cell culture applications at 10 mg/L $C_{22}H_{24}N_2O_8 \cdot HCl$ MW : 480.9 CAS : 64-75-5 Potency : $\geq 720 \text{ IU}/\text{mg}$ Store at 2 - 8°C	TC036-5G TC036-25G	5gm 25gm
Thimerosal Cell Culture Tested Thiomersal; Thiomersalate; Mercurothiolate; Sodium merthiolate; Sodium ethylmercurithiosalicylate $C_9H_9HgNaO_2S$ MW : 404.81 CAS : 54-64-8 Assay : $\geq 97\%$ Store below 30°C	TC470-10G TC470-25G TC470-100G TC470-500G	10gm 25gm 100gm 500gm
Tylosin tartrate Cell Culture Tested Potency : $\geq 900 \text{ U}/\text{mg}$ Store at 2 - 8°C	TC205-1G TC205-5G TC205-10G	1gm 5gm 10gm
Vancomycin hydrochloride Cell Culture Tested $C_{66}H_{75}Cl_2N_9O_{11} \cdot HCl$ MW : 1485.71 CAS : 1404-93-9 Assay : $\geq 80\%$ Potency : $\geq 900 \mu\text{g}/\text{mg}$ Store at 2 - 8°C	TC202-500MG TC202-1G TC202-5G	500mg 1gm 5gm
Gene Selection Antibiotics		
G418 disulfate salt (Geneticin disulphate) Cell Culture Tested Recommended for use in Cell Culture application at 100-800 mg/L $C_{10}H_{10}N_2O_{10} \cdot 2H_2SO_4$ MW : 692.71 CAS : 108321-42-2 Potency : $\geq 650 \mu\text{g}/\text{mg}$ Store at 2 - 8°C	TC025-1G TC025-5G	1gm 5gm
Hygromycin-B Cell Culture Tested Recommended for use in cell culture applications at 200-400 mg/L $C_{20}H_{37}N_3O_{13}$ MW : 527.52 CAS : 31282-04-9 Assay : $\geq 80\%$ Potency : $\geq 1170 \mu\text{g}/\text{mg}$ Store at 2 - 8°C	TC027-50MG TC027-100MG TC027-250MG TC027-1G	50mg 100mg 250mg 1gm

Product	Code	Packing
*Methotrexate Disodium Salt Cell Culture Tested $C_{20}H_{20}N_6Na_2O_5$ MW : 498.4 CAS : 7413-34-5 Assay : $\geq 98.5\%$ Store at 2 - 8°C	TC404-250MG	250mg
*Mycophenolic acid Cell Culture Tested $C_{20}H_{20}N_2Na_2O_5$ MW : 320.34 CAS : 24280-93-1 Assay : $\geq 99\%$ Store at 2 - 8°C	TC405-25MG TC405-50MG TC405-250MG	25mg 50mg 250mg
Neomycin sulfate salt Cell Culture Tested Recommended for use in cell culture application at 50 mg/L $C_{23}H_{46}N_6O_{13} \cdot 3H_2SO_4$ MW : 908.88 CAS : 1405-10-3 Potency : $\geq 600 \text{ IU}/\text{mg}$ Store below 30°C	TC031-5G TC031-25G	5gm 25gm
Puromycin dihydrochloride Cell Culture Tested $C_{22}H_{29}N_7O_5 \cdot 2HCl$ MW : 544.43 CAS : 58-58-2 Assay : $\geq 98\%$ Store at 2 - 8°C	TC198-10MG TC198-25MG	10mg 25mg
Anticancer Antibiotics		
*17-AAG 17-(Allylamino)-17-demethoxygeldanamycin Cell Culture Tested $C_{33}H_{43}N_3O_8$ MW : 585.69 CAS : 75747-14-7 Assay : $\geq 99\%$ Store at 2 - 8°C	TC409-1MG TC409-5MG	1mg 5mg
*Avermectin B1a, EvoPure® 5-O-Demethylavermectin A1a Cell Culture Tested $C_{48}H_{72}O_{14}$ MW : 873.1 CAS : 65195-55-3 Assay : $\geq 99\%$ Store at - 20°C	TC410-5MG	5mg
*Conglobatin (3Z,5R,7S,8S,11Z,13R,15S,16S)-3,5,7,11,13,15-hexamethyl-8,16-bis(1,3-oxazol-5-ylmethyl)-1,9-dioxacyclohexadeca-3,11-diene-2,10-dione Cell Culture Tested $C_{28}H_{38}N_2O_6$ MW : 498.61 CAS : 72263-05-9 Assay : $\geq 95\%$ Store at - 20°C	TC411-1MG	1mg
*Erlotinib Hydrochloride Cell Culture Tested $C_{22}H_{23}N_3O_4$ MW : 492.9 CAS : 183321-74-6 Assay : $\geq 98\%$ Store at 2 - 8°C	TC412-1G TC412-5G	1gm 5gm

*Delivery after 4 - 6 weeks

Unless indicated all the above products to be stored below 30°C.

Avermectin B1a, EvoPure® is a registered trademark of Toku-E company